

# NERVE BLOCKS IN ED TRUNK

陳國智醫師 雙和醫院急診醫學科

[juice119@gmail.com](mailto:juice119@gmail.com)

[POCUSacademy.com](http://POCUSacademy.com)

Resuscitative  
急救復甦

Diagnostic  
臨床診斷

Procedural  
Guidance  
處置導引

Symptom- or  
Sign-Based  
症狀導引

Therapeutic  
輔助治療



Core Applications (2023 ACEP Emergency Ultrasound Guidelines)  
15項急診超音波核心應用

陳國智醫師

Aorta

DVT

Trauma

Thoracic/Airway

Cardia/HD assessment

Procedural Guidance

US-guided NB

Testicular

Ocular

Skin & Soft tissue

Hepatobiliary

Urinary tract

Pregnancy

Bowel


MSK

Review

# The Pain Management of Trauma Patients in the Emergency Department


Andrea Fabbri <sup>1,\*</sup>, Antonio Voza <sup>2</sup>, Alessandro Riccardi <sup>3</sup>, Sossio Serra <sup>4</sup>  
and Fabio De Iaco <sup>5</sup> on behalf of the Study and Research Center of the Italian Society of Emergency Medicine (SIMEU)

The Multi-Modal Trauma Pain Approach



# NERVE BLOCK MANUAL

First Edition



Prof. Dr. Admir Hadzic

熟悉解剖


臨床情境

穩定運針

辨識血管

利弊分析

小心LAST



# Ultrasound for Interventional Pain Management





An Illustrated Procedural Guide





Philip Peng  
Roderick Finlayson  
Sang Hoon Lee  
Anuj Bhatia  
*Editors*

## The Ultimate Guide to Point-of-Care Ultrasound-Guided Procedures

Srikar Adhikari  
Michael Blaiwas  
*Editors*

**2024 ICEM**





# NIPRO

単回使用皮下注射用針

## ニプロカテラン針GA

管理医療機器  
ガンマ線照射済 使用1回限り

### 100本入

1. 包装及び製造過程中、用印止め表示漏れ。  
2. 露光存放時、照射面を直射日光にさらす。  
3. 照射面を損傷する。  
以上事項、発生、有害放射能汚染の責任を負担。

ニプロカテラン針GA  
ニプロ 21G x 70mm | R.B.

# NIPRO

単回使用皮下注射用針

## ニプロカテラン針GA

管理医療機器  
ガンマ線照射済 使用1回限り


### 100本入

1. 包装及び製造過程中、用印止め表示漏れ。  
2. 露光存放時、照射面を直射日光にさらす。  
3. 照射面を損傷する。



# SCP sensory distribution



The ultimate guide to POCUS-guided procedures



- Superficial cervical plexus
- Lesser occipital nerve
- Greater auricular nerve
- Transverse cervical nerve
- Supraclavicular nerve

# SCP block: in-plane approach


The ultimate guide to POCUS-guided procedures



LA: 5-8 ml

# SCP block

F 6.6~13.0  
DR 115  
FR 45  
D 2.5  
G 60



iNeedle

iTouch


thk: mm

DFOV:

KV:  
mséc  
ALC

# Supraclavicular nerve

DR 115  
FR 45  
D 2.5  
G 60



iNeedle

iTouch

thk: mm

DFOV:

KV:  
mséc  
ALG:

# Supraclavicular nerve




# SCN for clavicle fracture



# SCN for clavicle fracture

DR 115  
FR 45  
D 2.5  
G 73




iNeedle

iTouch

# SCN for clavicle fracture




F 6.6~13.0  
DR 115  
FR 45  
D 2.5  
G 70



iNeedle

iTouch

# Clavipectoral fascial plane block




# Regional anesthesia for clavicle fractures in emergency medicine: A scoping review


En-Hsien Su, Cheng-Chien Chen, Kuo-Chih Chen, Su Weng Chau, Yi-Kung Lee, Tou-Yuan Tsai ✉

First published: 26 December 2024 | <https://doi.org/10.1002/hkj2.12078>

<b>Sensory Innervation</b>	<b>Dermatomes</b>	<b>Myotomes</b>	<b>Osteotomes</b>
Involved root	C3–C5	C3–C7	C3–C7
Individual nerve	<ol style="list-style-type: none"><li>1. Supraclavicular nerve (C3–C4)</li><li>2. Ventral rami of C5</li></ol>	<ol style="list-style-type: none"><li>1. Axillary nerve (C5–C6)</li><li>2. Suprascapular nerve (C5–C6)</li><li>3. Lateral Pectoral nerve (C5–C7)</li><li>4. Nerve to subclavius (C5–C6)</li><li>5. Spinal accessory nerve (CN XI)</li><li>6. Trapezius branch of cervical nerves (C3–C4)</li></ol>	<ol style="list-style-type: none"><li>1. Axillary nerve (C5–C6)</li><li>2. Suprascapular nerve (C5–C6)</li><li>3. Lateral Pectoral nerve (C5–C7)</li><li>4. Supraclavicular nerve (C3–C4)</li><li>5. Nerve to subclavius (C5–C6)</li><li>6. Spinal accessory nerve (CN XI)</li><li>7. Trapezius branch of cervical nerves (C3–C4)</li></ol>

- Periosteal Collar
- Clavipectoral Fascia






SCPB and supraclavicular nerve block	Supine or semisitting position, with head turned away from the site being injected, exposing patient's SCM posterior border	Midpoint of SCM posterior border or midpoint of the mastoid process to C6 vertebra (Chassaignac tubercle)	5–15 mL local anesthetics	Greater auricular, lesser occipital, transverse cervical, and suprascapular nerves
Interscalene nerve blocks	Supine lateral decubitus position, with head turned away from the site being injected	Between the anterior and middle scalene muscles, 3–4 cm superior to the clavicle	7–15 mL of local anesthetic	Superior and middle trunks of brachial plexus
Clavipectoral plane block	Supine position with the head of the bed elevated	Between the clavipectoral fascia and the periosteum of the clavicle	15 mL of local anesthetic	Blocking nerve terminal branches pass through between the clavipectoral fascia and the clavicle


# Clavipectoral fascial plane block

F 6.6~13.0  
DR 115  
FR 45  
D 25  
G 70




iNeedle

iTouch



7/06/07  
4:15



# Intercostal Block

## ▶▶ QUICK FACTS

**Indications:** Analgesia for rib fractures, postsurgical analgesia for chest and upper abdominal surgery (i.e., thoracotomy, thoracostomy, mastectomy, gastrostomy, and cholecystectomy), herpes zoster or post-herpetic neuralgia.

**Goal:** Local anesthetic spread in the intercostal sulcus around the intercostal nerve.

**Patient position:** An intercostal nerve block can be performed with the patient in the seated, lateral decubitus, or prone positions.

**Landmarks:** The US transducer is placed parasagittal to identify the ribs.



Linear




22-25 gauge,  
3-5 cm



3-5 mL at each level




# T2-T12 spinal nerve - thoracic wall & upper abdomen




NYSORA


Fig-1. Anatomy of the thoracic spinal and intercostal nerves.



# Intercostal nerve block



# Intercostal nerve block



**Correct level**


**Inferior border**

**3-5ml/IC level**

**Hydrodissection**

**Consider 3 levels**

# Intercostal nerve block





U  
F-F5.0  
DR 10.0  
FR 71  
D 7.5  
G A 8

iNeedle



iTouch

LTZ 45 AP 96.5% MI 55 IS 0.1

U  
F-F5.0  
DR 10.0  
FR 71  
D 7.5  
G A 8

iNeedle

iTouch



# PVB (paravertebral block)

**Indications:** Analgesia after breast surgery, thoracotomy, rib fractures, and procedures involving the thoracic and upper abdominal wall.

**Goal:** Infiltration of local anesthetic into the paravertebral space.

**Patient position:** The PVB is frequently performed with the patient in a sitting position. Alternatively, it can be performed with the patient in the prone or in the lateral position with the side to be blocked facing upwards.

**Landmarks:** The transducer is placed parasagittal, about 3 cm from the midline, to identify the transverse processes.




Linear



21-22 gauge,  
8-10 cm





4-5 mL per space




# PVB (paravertebral block)


Superior costotransverse ligament  
Parietal pleura  
Vertebral body & intervertebral disk






# PVB (paravertebral block)






Identify dermatome



# PVB for postherpetic neuralgia



# Pectoralis / Serratus Plane block

**Indications:** PEC I: Analgesia for breast surgery, clavicle, and acromioclavicular procedures.

PEC II: Extensive breast surgery, thoracic wall procedures, sentinel node excision.

Serratus anterior plane: Anterolateral thoracic wall surgery and rib fractures.

**Goal:** Injection of local anesthetic in the fascial planes between the pectoral muscles and the serratus anterior muscle.

**Patient position:** Supine or semi-lateral position with the ipsilateral arm abducted 90 degrees.

**Landmarks:** The clavicle, pectoralis major muscle, anterior and midaxillary lines.



Linear



22 gauge,  
5-10 cm short bevel





10-20 mL







# Lateral and medial pectoral nerves


- Intercostobrachial nerve
- Pectoralis minor
- Thoracodorsal nerve
- Serratus anterior
- Pectoralis major (cut)




# PEC I & II block (pectoral fascial)




# SAPB (serratus anterior plane block)



# SAPB (serratus anterior plane block)






# SAPB (serratus anterior plane block)











20 ml  
Rib 2



10+20 ml  
Rib 4




20 ml  
Rib 5  
Mid-axillary  
Latissimus dorsi





# Coracobrachialis muscle abscess



CES:  
13.0  
25/03/04



19007BER

dle

ch  
: mm


19007BER

OV:



SE  
IM

K  
ms  
AL



SE: 1<sup>0</sup>  
IM: 15



-5  
KV:  
msec  
ALG:



# ESPB (erector spinae plane block)

**Indications:** Analgesia for rib fractures, back and chest wall surgeries.

**Goal:** Injection of local anesthetic in the plane deep to the erector spinae muscles and superficial to the transverse processes, to achieve a craniocaudal distribution along several vertebral levels.

**Patient position:** The patient is best positioned in a sitting, lateral decubitus or prone position. Identify the desired level for injection by counting down from the first rib (with ultrasound), or by taking into account the bony landmarks of the back, and their corresponding vertebral levels.

**Landmarks:** Midpoint of the clavicle and the coracoid process.




Linear or curved



22 - gauge,  
5-10 cm short bevel




20-30 mL



# ESPB (erector spinae plane block)


## Interfascial spreading




Spinalis

Longissimus

Iliocostalis




# ESPB (Interfascial spreading)




# ESP B (erector spinae plane block)




# ESPB (erector spinae plane block)






# ESPB (erector spinae plane block)






# ESPB (erector spinae plane block)






# ESPB (Rib level)




# ESPB (Rib / TP level)




# ESPB (lamina level)





# ESPBB (TP level - target of ESPBB)





# ESPB (TP level - target of ESPB)




# ESPB (TP level - target of ESPB)



# ESPB (TP level - target of ESPB)










# ESPB (TP level - target of ESPB)





# ESPB (TP level - target of ESPB)

Fascial block: at least 20 ml








# Sternal fracture hematoma block



# Sternal fracture hematoma block





P  
R  
1.1 8.8




Article

# Ultrasound Guided Parasternal Block for Perioperative Analgesia in Cardiac Surgery: A Prospective Study


Giuseppe Pascarella <sup>1</sup> , Fabio Costa <sup>1</sup>, Giulia Nonnis <sup>2</sup>, Alessandro Strumia <sup>1,\*</sup> , Domenico Sarubbi <sup>1</sup>, Lorenzo Schiavoni <sup>1</sup> , Annalaura Di Pumpo <sup>1</sup>, Lara Mortini <sup>1</sup>, Stefania Grande <sup>1</sup>, Andrea Attanasio <sup>3</sup>, Giovanni Gadotti <sup>4</sup>, Alessandro De Cassai <sup>5</sup> , Alessia Mattei <sup>1</sup>, Antonio Nenna <sup>6</sup> , Massimo Chello <sup>6</sup> , Rita Cataldo <sup>1</sup>, Felice Eugenio Agrò <sup>1</sup> and Massimiliano Carassiti <sup>1</sup>




# Parasternal block for Sternal fracture





# Lumbar ESPB




# Lumbar ESPB - parasagittal




# Lumbar ESPB - transverse




# Lumbar ESPB - Aksu approach




# Lumbar ESPB





# Lumbar ESPB




# Lumbar ESPB




# Lumbar ESPB






# Lumbar ESPB




# ESPB - L3 bursae fx



# Lumbar ESPB





# TAP Block

**Indications:** Analgesia for the abdominal wall and the parietal peritoneum.


**Goal:** Local anesthetic spread in the transversus abdominis plane (TAP): The plane between the internal oblique and transversus abdominis muscles.

**Patient position:**

- For the lateral and subcostal TAP blocks: Supine.
- For the posterior TAP block: Lateral decubitus.

**Landmarks:**

- **Subcostal TAP block:** Subcostal margin and xiphoid process.
- **Lateral and posterior TAP blocks:** Iliac crest and midaxillary line.
- **Ilioinguinal-iliohypogastric block:** Anterior superior iliac spine and the umbilicus.



Linear or curved  
(larger patients)



22 gauge,  
5-10 cm



15-30 mL



Dorsal root  
(Sensory root)

Ventral root  
(Motor root)

Spinal ganglion

Meningeal ramus

Spinal nerve

Dorsal ramus (posterior)  
with medial ramus and  
lateral ramus


Ramus communicans


Ventral ramus

Sympathetic ganglion

Lateral cutaneous ramus

Ventral cutaneous ramus









*Fig-3. Expected sensory distribution of the different approaches for a TAP block.*

# TAP Block - surface landmark


triangle of Petit



# Transversus Abdominis Plane Blocks



USG subcostal TAP block



USG lateral TAP block

Lateral cutaneous branches of thoracoabdominal nn. (T6-T12)


Anterior cutaneous branches of thoracoabdominal nn. (T6-T12)

External oblique m. and aponeurosis


# TAP Block



# Anterior TAP (Ilioinguinal-iliohypogastric)




# Anterior TAP (Ilioinguinal-iliohypogastric)



# Anterior TAP (Ilioinguinal-iliohypogastric)


Arterial  
L12-3  
49 Hz  
2.5cm

2D  
HGen  
Gn 66  
C. 41  
3/3/2










1.1 8.8

# Anterior TAP (Ilioinguinal-iliohypogastric)




P  
R  
1.8 3.6



7.0cm






# Rectus sheath block



NYSORA®



# Rectus sheath block



# Quadratus lumborum block

**Indications:** Analgesia for the anterolateral abdominal wall and parietal peritoneum.

**Goal:** Local anesthetic injection lateral (QL1), posterior (QL2) or anterior (QL3) to the quadratus lumborum muscle.

**Patient position:**

- The lateral decubitus position is preferred over the supine position as it provides better ergonomics and a better view of the relevant ultrasound images and neuraxis. The QL1, QL2 and QL3 blocks can be performed in this position.
- The supine position is suitable only for the QL1 approach.

**Landmarks:** Iliac crest, costal margin, and the posterior and midaxillary lines.



Linear or Curved




22 gauge,  
5-10 cm short bevel



15-20 mL



# TLF (thoracolumbar fascia)



# Quadratus lumborum (T7-L1)



Aortic opening

Lateral arcuate ligament

Medial arcuate ligament

12th rib


Quadratus lumborum muscle

Psoas muscle


Iliacus muscle

# QL block



(T7-12 / Paravertebral space / sympathetic chain)





# QL 1 (lateral) block



# QL 2 (posterior) block






# QL 3 (anterior) block






**Shamrock sign**


# Caudal block




# Caudal block





# Caudal block



# Caudal block





# Caudal block for chronic low back pain






# Caudal block for radiculopathy



# Caudal block for radiculopathy




# Take Home Skills



Hospital MSK 3 3158 10:45:11 AM

MSK FR TIS DL2 5:13:12 PM



Depth Freeze Save Optic

# TRUNK BLOCKS

<b>Clavipectoral block</b>	<b>Intercostal block</b>	<b>Rectus sheath block</b>
<b>Parasternal block</b>	<b>PVB</b>	<b>TAP - subcostal</b>
<b>PEC I</b>	<b>ESPB - thoracic</b>	<b>TAP - lat / posterior</b>
<b>PEC II</b>	<b>ESPB - lumbar</b>	<b>TAP - ant (IN-IH)</b>
<b>SAPB</b>	<b>Caudal block</b>	<b>QL block (1 / 2 / 3 )</b>