



PICC


Peripherally Inserted Central Catheter

陳國智醫師 雙和醫院急診醫學科


juice119@gmail.com / POCUSacademy.com

先進與革新急症技術中心 / Advanced & Revolutionary Technology center

A. Peripheral IV Catheter




D. Nontunneled Central Venous Catheter




The Michigan Appropriateness Guide for Intravenous Catheters (MAGIC): Results From a Multispecialty Panel Using the RAND/UCLA Appropriateness Method


Vineet Chopra, MD, MSc; Scott A. Flanders, MD; Sarjay Saint, MD, MPH; Scott C. Wolter, MD; Naomi P. O'Grady, MD; Nasir Sadiq, MD, PhD; Scott O. Trerotola, MD; Rajiv Saren, MD, PhD; Nancy Moreau, BSN, RN; Stephen Wiseman, PharmD; Mauro Piktiri, MD; Ellie A. Akl, MD, MPH, PhD; Agnes Y. Lee, MD, MSc; Anthony Courcy, MD; Lakshmi Swaminathan, MD; Jack LeDonne, MD; Carol Becker, MHA; Sarah L. Krein, PhD, RN; and Steven J. Bernstein, MD, MPH




B. US-Guided Peripheral IV Catheter




G. Peripherally Inserted Central Catheter




E. Tunneled Central Venous Catheter



C. Midline Catheter



F. Implanted Port



週邊置入中心靜脈導管

**G. Peripherally Inserted
Central Catheter**



MAGIC

Michigan Appropriateness Guide
for Intravenous Catheters



經週邊給予輸液時的建議

Chopra V, Ann Intern Med, 2015

Device Type	Proposed Duration of Infusion			
	≤5 d	6–14 d	15–30 d	≥31 d
Peripheral IV catheter	No preference between peripheral IV and US-guided peripheral IV catheters for use ≤5 d			
US-guided peripheral IV catheter	US-guided peripheral IV catheter preferred to peripheral IV catheter if proposed duration is 6–14 d			
Nontunneled/acute central venous catheter	Central venous catheter preferred in critically ill patients or if hemodynamic monitoring is needed for 6–14 d			
Midline catheter	Midline catheter preferred to PICC if proposed duration is ≤14 d			
PICC		PICC preferred to midline catheter if proposed duration of infusion is ≥15 d		
Tunneled catheter				PICC preferred to tunneled catheter and ports for infusion 15–30 d
Port				

Appropriate

Neutral

Inappropriate

Disagreement

無法由週邊管路治療時的建議

Chopra V, Ann Intern Med, 2015

Device Type	Proposed Duration of Infusion			
	≤5 d	6–14 d	15–30 d	≥31 d
Peripheral IV catheter				
US-guided peripheral IV catheter				
Nontunneled/acute central venous catheter	Central venous catheter preferred in critically ill patients or if hemodynamic monitoring is needed for 6–14 d			
Midline catheter				
PICC		PICCs rated as appropriate at all proposed durations of infusion		
Tunneled catheter		Tunneled catheter neutral for use ≥15 d	No preference between tunneled catheter and PICC for proposed durations ≥15 d	
Port				No preference among port, tunneled catheter, or PICC for ≥31 d

5

Appropriate

Neutral

Inappropriate

Disagreement

困難靜脈管路建立時的建議

Chopra V, Ann Intern Med, 2015

Device Type	Proposed Duration of Infusion			
	≤5 d	6–14 d	15–30 d	≥31 d
Peripheral IV catheter	No preference between peripheral IV and US-guided peripheral IV catheters for use ≤5 d			
US-guided peripheral IV catheter	US-guided peripheral IV catheters preferred to peripheral IV catheters if proposed duration is 6–14 d			
Midline catheter	Midline catheters preferred to PICC if proposed duration is ≤14 d			
Nontunneled/acute central venous catheter	Central venous catheter preferred to PICC for use ≤14 d in critically ill patients			
PICC	Disagreement on appropriateness of PICC for durations <5 d	PICC use appropriate if proposed duration is ≥6 d; PICCs preferred to tunneled catheters for durations of 15–30 d		
Tunneled catheter			Tunneled catheter neutral for difficult IV access for use ≥15 d	No preference between tunneled catheter or port for use ≥31 d
Port				

6

Appropriate

Neutral

Inappropriate

Disagreement

需要經常抽血時的建議

Chopra V, Ann Intern Med, 2015

Device Type	Proposed Duration of Infusion			
	≤5 d	6–14 d	15–30 d	≥31 d
Peripheral IV catheter	No preference between peripheral IV and US-guided peripheral IV catheter for use ≤5 d US-guided peripheral IV catheter preferred if venous access difficult			
US-guided peripheral IV catheter				
Midline catheter	Midline catheter preferred to PICCs if proposed duration is ≤14 d		Midline catheter neutral for frequent phlebotomy at this duration	
Nontunneled/acute central venous catheter	Central venous catheter preferred to PICC for use ≤14 d in critically ill patients			
PICC	Disagreement on appropriateness of PICC for durations <5 d	PICC use appropriate if proposed duration ≥6 d; PICC preferred to tunneled catheter for durations of 15–30 d		
Tunneled catheter			Tunneled catheter neutral for difficult intravenous access for use ≥15 d	
Port	Ports inappropriate for frequent phlebotomy, regardless of proposed duration of use			

Appropriate

Neutral

Inappropriate

Disagreement

Which of the following indication are you ordering a PICC for?

(select one)

☐ IV Antibiotics

☐ Vesicant or Irritant Chemotherapy

☐ Non chemotherapy Vesicant or Irritant

☐ Difficult Venous Access

☐ Frequent Blood Draws

What is your proposed duration of therapy?

(select one)

☐ Less than 5 days

☐ 6-14 days

☐ 15-30 days

☐ More than 31 days

Appropriate

Inappropriate

Click (i) for preference recommendations

PICC

Tunneled Catheter

Peripheral IV Catheter

Ultrasound Guided Peripheral Catheter

Non-tunneled/Acute Central Venous Catheter



PICCs rated as appropriate at all proposed duration of Infusion

Click here to see the MAGIC algorithm

OK

Critical and Honest Conversations

The Evidence Behind the “Choosing Wisely” Campaign Recommendations by the American Society of Nephrology

Williams, Amy W.; Dwyer, Amy C.; Eddy, Allison A.; Fink, Jeffrey C.; Jaber, Bertrand L.; Linas, Stuart L.; Michael, Beckie; O'Hare, Ann M.; Schaefer, Heidi M.; Shaffer, Rachel N.; Trachtman, Howard; Weiner, Daniel E.; Falk, and Ronald J.

[Author Information](#) ✓

Clinical Journal of the American Society of Nephrology 7(10):p 1664-1672, October 2012. | DOI: 10.2215/CJN.04970512

Avoid PICC on Patients with eGFR <45ml/min

4. Preserving veins for future creation of an arteriovenous fistula (AVF) is paramount to ensure better outcomes for patients with CKD who may eventually need HD.

CWCr: “Do not place peripherally inserted central venous catheters (PICCs) in stage 3–5 CKD patients without consulting nephrology.”





Peripherally Inserted Central Catheter lines for Intensive Care Unit and onco-hematologic patients: A systematic review and meta-analysis

Georgios Mavrovounis, MD^{a,*}, Maria Mermiri, MD^a, Dimitrios G Chatzis, MD, PhD^b, Ioannis Pantazopoulos, MD, MSc, PhD^a

^a Department of Emergency Medicine, Faculty of Medicine, School of Health Sciences, University of Thessaly, General University Hospital of Larissa, Mezourlo 41110, Larissa, Greece

^b School of Medicine, European University of Cyprus, Nicosia, Cyprus, Diogenous Str 2404, Nicosia, Cyprus

Mavrovounis G, et al. Heart Lung 2020



Thrombosis risk
PICC > CVC

CLABSI risk
CVC > PICC onco-hematologic patients

PICC併發症

Infection (CRBSI 1.1/1000 PICC days)

Catheter malposition / migration

Mechanical malfunction (10 ~ 27%)

Phlebitis / infiltration (2.2% ~ 23%)

Air embolism

Cardiac arrhythmia

Catheter occlusion-thrombotic

High SVC 78% v.s. Distal SVC/RA 16%

Catheter occlusion-non-thrombotic

常用的靜脈導管比較

黃耀廣醫師網站 [HTTPS://WWW.CVSDOCTOR.COM/PICC-INTRODUCTION/](https://www.cvdoctor.com/picc-introduction/)

輸液導管	週邊靜脈留置針	週邊置入中心靜脈導管	中央靜脈導管
留置時間	需要每3天更換	半年以上 😊	每7至10 天需更換
置放者	由護理師置放	由醫師置放	由醫師置放
功能	當靜脈輸液的酸鹼值和滲透壓值和血液不同時，容易刺激血管內皮細胞而造成損傷，進而形成化學性的血栓	可監測中心靜脈壓(威力週邊置入中心導管、中央靜脈導管) 有雙腔、三腔管路，可同時提供病人多種或大量輸液 具刺激性之化學治療藥物建議應由中心靜脈導管給予，尤其是起皰性的化學治療藥物，為了避免藥物外滲時所造成的重大傷害	
合併症	無重大併發症 可能引起靜脈炎、藥物外滲、皮膚壞死	無重大併發症 😊	放置過程可能會產生氣、血胸、大動脈穿刺、血栓感染機率較高
抽血	每次抽血均需重覆扎針	放置後不需重覆紮針，自外露管路抽血即可	放置後不需重覆紮針，自外露管路抽血即可
電腦斷層	由留置針施打顯影劑可能會造成顯影劑外滲	自管路直接施打顯影劑即可(威力週邊置入中心導管) 😊	仍需另放置週邊靜脈留置針(可能會造成顯影劑外滲)
放置部位	前臂或手易影響活動	手臂上段內側，不會影響活動 😊	頸部或鼠蹊部，易影響活動

PICC在病人端的好處

不會影響活動、較舒適和美觀

抽血給葯方便、不必反覆挨針

葯物注射時無疼痛、化療葯物不外漏較安全

可以保護血管、療程結束後即可移除

PICC在醫療端的好處

低侵入性、重大合併症不易發生

可長期放置、可給特殊性輸液及葯物

臨床照護較CVC容易、感染率也較低

PRESSURE INJECTABLE (CT)

第六卷第四期

刊登日期：2023/08/31

Taiwan Emergency Medicine Bulletin 6(4) : e2023060408

How I do it: 如何在急診開始第一隻PICC？

下載PDF

LINE 分享

Facebook 分享

王鎮琮、范閔皓、陳國智 、翁健瑞

衛生福利部雙和醫院 急診醫學科

台灣急診醫學通訊



第六卷第四期

刊登日期：2023/08/31

Taiwan Emergency Medicine Bulletin 6(4) : e2023060408

How I do it: 如何在急診開始第一隻PICC？

下載PDF

LINE 分享

Facebook 分享


王鎮琮、范閔皓、陳國智 、翁健瑞

衛生福利部雙和醫院 急診醫學科

台灣急診醫學通訊



PICC 週邊置入中心靜脈導管



適應情境

置放選擇

導管置放

確認位置

管路照護

Teleflex Pressure Injectable PICC

1. BlueFlex Tip®

Soft, reduces damage to delicate intima during threading process¹

2. Staggered Exit Ports

Reduce risk of mixing incompatible drugs and solutions that may create precipitate²

3. Labeled Depth Marker


Every 1 cm on Catheter Body

4. Increase Radiopacity of Catheter Body

6. Indicate Maximum Injection Rates

5. Taper-Free Catheter

Allow for increase blood flow around the catheter and minimizes risk of Catheter related thrombosis



PICC健保支付標準 - 4適應症

47065B 點數 **3101** (111.3.1 ~)


癌症化學治療及癌症末期之疼痛治療

下列三項適應症且預期同時治療達**二週以上**之病人

- 使用全靜脈營養輸液(TPN)(1歲以下嬰兒本項為靜脈營養輸液，不限TPN)
- 免疫不全與使用免疫抑制劑
- 須接受中心靜脈導管置入(CVC)



PICC可放置期間



6個月 ~ 1年

The SLP protocol update: Eight strategies, incorporating Rapid Peripheral Vein Assessment (RaPeVA), to minimize complications associated with peripherally inserted central catheter insertion

Fabrizio Brescia¹ , Mauro Pittiruti² ,
Timothy R Spencer³  and Robert B Dawson⁴

The Journal of Vascular Access

1–9

© The Author(s) 2022



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/11297298221099838

journals.sagepub.com/home/jva



SIP PROTOCOL


Safe insertion of PICCs

- Step 1 *Pre-procedural evaluation*—choose most appropriate vein by systematic ultrasound examination of the veins of the arms (see the RaPeVA protocol)
- Step 2 *Appropriate antiseptic technique*—adopt a strict policy of hand hygiene, skin antisepsis with 2% chlorhexidine in 70% isopropyl alcohol, and use of maximal barrier precautions
- Step 3 *Choice of vein size and exit site*—evaluate the diameter of the vein so to have an ideal catheter-vein ratio (1:3 or less); place the exit site in the green zone (see Dawson's ZIM™); consider the opportunity of tunneling the catheter, if the most appropriate vein is in the yellow zone (see the RAVESTO protocol)
- Step 4 *Clear identification of median nerve and brachial artery*—identify each structure before venipuncture, using ultrasound
- Step 5 *Ultrasound-guided venipuncture*—access a deep vein of the arm (either basilic or brachial vein), preferably adopting the short axis/out-of-plane approach, and use of a micro-introducer kit
- Step 6 *Ultrasound-based tip navigation*—assess the correct direction of the guidewire, by a supra-clavicular ultrasound scan (see the ECHOTIP protocol)
- Step 7 *Intra-procedural assessment of tip location*—use intracavitary ECG and/or ultrasound (subcostal or apical view, using the “bubble test”: see the ECHOTIP protocol)
- Step 8 *Appropriate securement of the catheter and protection of the exit site*—use sutureless devices only; reduce the risk of bleeding and bacterial contamination using cyanoacrylate glue and semi-permeable transparent membrane dressings

RaPeVa protocol

Rapid Peripheral Vein Assessment


- | | |
|--------|---|
| Step 1 | Visualization of the cephalic vein at the antecubital fossa |
| Step 2 | Identification of the artery and brachial veins and of the confluence between the antecubital vein and basilic vein |
| Step 3 | Identification of the basilic vein in the bicipital-humeral groove |
| Step 4 | Examination of the nerve-vascular bundle of the arm |
| Step 5 | Visualization of the cephalic vein over the biceps muscle |
| Step 6 | Examination of the axillary vein in the infraclavicular area |
| Step 7 | Examination of the internal jugular, the subclavian, and the brachio-cephalic vein in the supraclavicular area |



RaPeVa protocol

Rapid Peripheral Vein Assessment


- | | |
|--------|---|
| Step 1 | Visualization of the cephalic vein at the antecubital fossa |
| Step 2 | Identification of the artery and brachial veins and of the confluence between the antecubital vein and basilic vein |
| Step 3 | Identification of the basilic vein in the bicipital-humeral groove |
| Step 4 | Examination of the nerve-vascular bundle of the arm |
| Step 5 | Visualization of the cephalic vein over the biceps muscle |
| Step 6 | Examination of the axillary vein in the infraclavicular area |
| Step 7 | Examination of the internal jugular, the subclavian, and the brachio-cephalic vein in the supraclavicular area |



RaPeVa protocol

Rapid Peripheral Vein Assessment


- | | |
|--------|---|
| Step 1 | Visualization of the cephalic vein at the antecubital fossa |
| Step 2 | Identification of the artery and brachial veins and of the confluence between the antecubital vein and basilic vein |
| Step 3 | Identification of the basilic vein in the bicipital-humeral groove |
| Step 4 | Examination of the nerve-vascular bundle of the arm |
| Step 5 | Visualization of the cephalic vein over the biceps muscle |
| Step 6 | Examination of the axillary vein in the infraclavicular area |
| Step 7 | Examination of the internal jugular, the subclavian, and the brachio-cephalic vein in the supraclavicular area |



RaPeVa protocol

Rapid Peripheral Vein Assessment

- | | |
|--------|---|
| Step 1 | Visualization of the cephalic vein at the antecubital fossa |
| Step 2 | Identification of the artery and brachial veins and of the confluence between the antecubital vein and basilic vein |
| Step 3 | Identification of the basilic vein in the bicipital-humeral groove |
| Step 4 | Examination of the nerve-vascular bundle of the arm |
| Step 5 | Visualization of the cephalic vein over the biceps muscle |
| Step 6 | Examination of the axillary vein in the infraclavicular area |
| Step 7 | Examination of the internal jugular, the subclavian, and the brachio-cephalic vein in the supraclavicular area |



RaPeVa protocol

Rapid Peripheral Vein Assessment


- | | |
|--------|---|
| Step 1 | Visualization of the cephalic vein at the antecubital fossa |
| Step 2 | Identification of the artery and brachial veins and of the confluence between the antecubital vein and basilic vein |
| Step 3 | Identification of the basilic vein in the bicipital-humeral groove |
| Step 4 | Examination of the nerve-vascular bundle of the arm |
| Step 5 | Visualization of the cephalic vein over the biceps muscle |
| Step 6 | Examination of the axillary vein in the infraclavicular area |
| Step 7 | Examination of the internal jugular, the subclavian, and the brachio-cephalic vein in the supraclavicular area |



RaPeVa protocol

Rapid Peripheral Vein Assessment



- | | |
|--------|---|
| Step 1 | Visualization of the cephalic vein at the antecubital fossa |
| Step 2 | Identification of the artery and brachial veins and of the confluence between the antecubital vein and basilic vein |
| Step 3 | Identification of the basilic vein in the bicipital-humeral groove |
| Step 4 | Examination of the nerve-vascular bundle of the arm |
| Step 5 | Visualization of the cephalic vein over the biceps muscle |
| Step 6 | Examination of the axillary vein in the infraclavicular area |
| Step 7 | Examination of the internal jugular, the subclavian, and the brachio-cephalic vein in the supraclavicular area |




RaPeVa protocol

Rapid Peripheral Vein Assessment

- Step 1 Visualization of the cephalic vein at the antecubital fossa
- Step 2 Identification of the artery and brachial veins and of the confluence between the antecubital vein and basilic vein
- Step 3 Identification of the basilic vein in the bicipital-humeral groove
- Step 4 Examination of the nerve-vascular bundle of the arm
- Step 5 Visualization of the cephalic vein over the biceps muscle
- Step 6 Examination of the axillary vein in the infraclavicular area
- Step 7 Examination of the internal jugular, the subclavian, and the brachio-cephalic vein in the supraclavicular area



PICC置放選擇



APPROPRIATE VEIN ZON INSERTION METHOD (ZIM)

		Radius of Vessel (mm) ⁴	Length (CM)	Actual Diameter
Cephalic		3 ⁴	38 cm	2-4 mm
Basilic		4 ⁴	24 cm	4-6 mm
Axillary		8 ⁴	13 cm	16 mm
Subclavian		9.5 ⁴	6 cm	19 mm
SVC		12.5 ⁴	7 cm	20 mm

ZIM FOR EXIT SITE

Length of the arm
(Acromion - Olecranon)

Yellow: proximal third

Green: middle third

Red: distal third

C-V RATIO

1 : 3

1Fr 0.33mm

3Fr 1mm

US GUIDED PIV

Ⓐ

Landmark: humerus & brachial aretery

basilic v

V

A


V

brachial v

humerus

内側


外側





Brachial artery / vein & Basilic vein


BASILIC VEIN & BRACHIAL VEIN



CEPHALIC VEIN

外側

内側



MI
1.5
11L4
diffT9.0
30 fps
Qscan
G:81
DR:60
A:6
P:3

The 3 Stations of the Needle: A systematic approach for ultrasound guided venous and arterial puncture

Matthew Ostroff¹  and Nancy Moureau^{2,3} 

The Journal of Vascular Access
1–7

© The Author(s) 2025

Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/11297298251334888

journals.sagepub.com/home/jva






Off - plane



In - plane

Superficial P
L12-3
46 Hz
3.0cm
2D
Res
Gn 96
C 56
3/2/1



Superficial P
L12-3
46 Hz
3.0cm
2D
Res
Gn 96
C 56
3/2/1














Table 1. 3-Stations of the Needle.

Stations of the needle	Transverse (Short axis)	Longitudinal (Long axis)	Description	Key points
1 浅入針			Identification of the needle tip upon skin entry to the subcutaneous tissue	Procedural time out confirming needle trajectory and depth of vessel before advancement of the needle
2 抬針尾			Navigation of needle tip from the subcutaneous tissue to outer wall of target vessel	Procedural time out for assessment of the surrounding structures of the intended target vessel for safe needle navigation using the PN-T
3 找針尖			Vessel confirmation (arterial or venous) and needle advancement through vessel walls	Procedural time out before vessel puncture and P-NT for vessel purchase



斜面由朝上轉朝下


斜面由朝下轉朝上



送Guidewire斜面朝下

送Catheter則斜面朝上

OFF-PLANE APPROACH PREFERRED CHOICE FOR PICC








1




3



2



4




Optimal skin surface landmark for the SVC-RA junction in cancer patients requiring the implantation of permanent central venous catheters

J. H. Hsu,¹ S. S. Wang,² D. V. Lu,³ K. I. Cheng,³ C. K. Wang⁴ and J. R. Wu^{1,5}

Departments of 1 Paediatrics and 4 Medical Imaging, Kaohsiung Medical University Hospital, 100 Tzyou 1st Road, Kaohsiung 807, Taiwan

2 Department of Surgery, Chang Gung Memorial Hospital, no. 6, Section west, Chia Pu Road, Puzi City, Chiayi 613, Taiwan

Department of 3 Anaesthesiology and 5 Paediatrics, Kaohsiung Municipal Hsiao-Kang Hospital, 482 San-Ming Road,



CAJ: Lower margin of R 3rd

Table 3 Number and percentage of patients with satisfactory positioning guided by different skin surface landmarks (*n* = 20).

Skin surface landmark	No. of patients with satisfactory positioning	Percentage of patients with satisfactory positioning	p value*
Point A	0	0%	< 0.0001
Point B	4	20%	
Point C	14	70%	
Point D	6	30%	

PICC導管置放



PICC導管置放



TE7 ACE

LI2-4s

20230214-083501-4868

65Years

Vascular

14-02-2023

07:55:39

20230214-083501-4868

65Years

Vascular

14-02-2023

08:03:32

20230214-083501-4868

65Years

Vascular




14-02-2023

08:05:40

B
F100.0
DR 95
FR 40
D 2.5
G 64

Needle

Touch






US-GUIDED PICC TIP NAVIGATION

B
F100.0
DR 95
FR 40
D 4.0
G 64

m

Needle

Touch



TE7 ACE

20230214-083501-486B

Vascular

14-02-2023
08:15:47

L12-4s

AP 96.6%

MI135


TIS 0.2

B
FH10.0
DR 95
FR 40
D 4.0
G 64

m

iNeedle

iTouch



L12-4s

AP 96.6%

MI135


TIS 0.2

portable R

20230214-083501-486B

Vascular

14-02-2023
08:20:50



-0

-1

-2

-3

ECHOTIP PROTOCOL TIP NAVIGATION

20230214-083501-486B Vascular 08:21:28

L12-4s

AP 95.6%

MI135


TIS 0.2

B
FH10.0
DR 95
FR 40
D 4.0
G 64

m

iNeedle

iTouch



20230214-083501-486B Vascular 08:21:38

AP 95.6%


MI135

TIS 0.2

m

iNeedle

iTouch



20230214-083501-486B Vascular 08:21:54

AP 95.6%


MI135

TIS 0.2

m

iNeedle

iTouch




-0


-1

-2




-3








TIP LOCALIZATION



PICC確認位置 (ECHO + CXR)









IM: 1
SE: 3
FH9.0
DB 100
PR 51
D 3.0
G 57
iNeedle
iTouch
thk: mm
PF19007BER
DFOV:




ChenKC@POCUSacademy




ChenKC@POCUSacademy








ChenKC@POCUSacademy



ChenKC@POCUSacademy



ChenKC@POCUSacademy



沖管



Q8H以10 mL N/S推停推停方式沖洗一次，

推停推停方式沖洗方式:快速推進1-2ml &每次推進
間隔約0.4秒¹



每次輸液前後、抽/輸血前後、更換管路及 / 或脂質
輸注時



沖洗必須涵蓋所有導管管腔

1. ISBN 978-3-030-03148-0 ISBN 978-3-030-03149-7 (eBook) <https://doi.org/10.1007/978-3-030-03149-7.S82>

採血/給予藥物

- 1.停止全身靜脈輸注液體或血液製品 2 分鐘
- 2.消毒無針連接器或導管銜接孔 **5-15秒**
- 3.以 10 mL N/S 推停推停方式沖洗管路
- 4.棄血 5 mL
- 5.連接新的10 mL 針筒
- 6.採血
- 7.消毒無針連接器銜接孔或導管銜接孔 **5-15秒** (使用適當消毒劑 如chlorhexidine, povidone iodine, 或70% alcohol，請務必注意，若後續換藥施加敷料覆蓋前，應待消毒液體完全乾燥)
- 8.採血後以 20 mL N/S 脈動式沖洗管路，及適當清潔導管及任何附加裝置

閉管

1. 停止使用時及施打藥物並完成最後沖洗之後，務必進行**正壓閉管**，保留0.5ml NS(勿全部推注完)以大拇指壓住針筒活塞後，接著關閉閥夾後再移除注射器，避免血液回流入導管內¹

1. N/S使用，請依醫院程序及注意事項

文獻出處:

1. Gorski LA, Hadaway L, Hagle ME, et al. Infusion therapy standards of practice. J Infus Nurs. 2021;44(suppl 1):S1-S224.
doi:10.1097/NAN.0000000000000396


PICC管路照護



PICC照護

A graphic featuring three concentric circles in yellow, orange, and red. In the center, the text "PICC照護" is written in a stylized, 3D font with a pink-to-white gradient and a shadow effect.

Avoid PICC on Patients with eGFR <45ml/min



MAGIC
Michigan Appropriateness Guide
for Intravenous Catheters